

Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>

available at [www.sciencedirect.com](http://www.sciencedirect.com)journal homepage: [www.elsevier.com/jbmt](http://www.elsevier.com/jbmt)

## SELF-MANAGEMENT: PATIENT SECTION

## Hip muscle training

D.C. Craig Liebenson\*

*L.A. Sports & Spine, 10474 Santa Monica Blvd, #304, Los Angeles 90025, United States*

Received 29 December 2010; accepted 29 December 2010

Low back and knee pain often are due to poor function of the hip joint. The hip is a large, deep ball and socket joint located near your groin. The outer gluteal muscles – the gluteus medius in particular – control the alignment of your knee and your pelvis during movement such as walking. For instance, during 85% of walking you are balancing on one

leg, and the gluteus medius muscle, and one side and lateral trunk muscles (quadratus lumborum) on the opposite side, are controlling your posture.

Isolated training of the gluteus medius muscle can be a first step in stabilizing a knee, hip, or back problem.

## The clam shell

- Lie on your side with knees bent;
- Place one hand on your pelvis (Figure 1a);
- Use a light band around your knees for resistance;




Figure 1 Clam shell.




Figure 2 Athletic position.

\* Tel.: +1 31047 02909; fax: +1 31047 03286.  
E-mail address: [craigliebenson@gmail.com](mailto:craigliebenson@gmail.com).




Figure 3 Functional clam shell.

- Raise your knee slowly while keeping your foot down (Figure 1b);
- Monitor with your hand that your pelvis & spine don't rotate;
- Perform 10–12 repetitions.

Once the gluteus medius muscle is “turned on” then it's activity should be integrated into functional activities.

### Functional clam shell

- Place a band around your knees;
- Stand in an athletic position (Figure 2);
- Allow your knee to slowly move inwards slightly (Figure 3a);
- Then, perform a functional clam shell by turning your knee out against the resistance of the band (Figure 3b);
- Perform 8–10 repetitions.

### The lateral band walk

- Start in an athletic position with a band just above your knees (Figure 4a);
- Balance on one leg (Figure 4b);
- Step to the side and balance again on the same leg (Figure 4c);
- Repeat for 5–6 steps;
- Reverse directions;
- Perform 2–3 repetitions;
- *Troubleshooting:* be sure to feel the buttocks or outer hip muscles on the leg you are balancing on & pushing off with;
  - If you feel the front of your thigh then “peel back” to the clam shell to wake up the hip/buttock muscles;
  - If you are feeling the lead leg that you are pulling with, then balance for longer on the support/push leg, and possibly perform a few shallow single leg squats on that leg;
  - To progress this exercise you may add a 2nd band around the ankles.




Figure 4 Lateral band walk.